

FluMist[®]

Influenza Virus Vaccine Live, Intranasal

Safety Monitoring

in Children Aged <5 Years

ADVISORY COMMITTEE ON IMMUNIZATION PRACTICES
Atlanta, GA
June 28, 2007

Karen R. Broder, MD

Lieutenant Commander, US Public Health Service
Senior Medical Advisor
Immunization Safety Office
Office of the Chief Science Officer
Centers for Disease Control and Prevention



General Approaches to Safety Monitoring for New Vaccines or Vaccines with New Indications

- Review pre-licensure safety data
- Identification of potential risks from phase 3 pre-licensure trials
- Review of available postmarketing data
- Vaccine Adverse Event Reporting System (VAERS) monitoring plan
- Vaccine Safety Datalink (VSD) project plan: assessment of risk through near real-time or planned studies
- Identification/creation of
 - Key case definitions
 - Candidate protocols
 - Need for special studies




Pre-licensure Primary Safety Study MICP11: Methods

- Multi-center, randomized (49% US)
 - Study group: FluMist[®]* + IM placebo (saline) (N=4243)
 - Active control group: trivalent inactivated influenza vaccine (TIV) + intranasal placebo (saline) (N=4232)
 - Subjects excluded if history of severe asthma or medically diagnosed or treated wheezing within 42 days before enrollment
- Two pre-specified safety wheezing outcomes from dose 1 through 42 days after the last vaccine dose
 - Primary: Medically significant wheeze
 - Secondary: Any wheeze



*Influenza Virus Vaccine, Trivalent types A and B, Live Cold Adapted, liquid formulation (2004-2005 influenza strains)

Sources: FDA and MedImmune briefing documents, FDA VRPBAC meeting, May 16, 2007; Belshe NEJM 2007.



Pre-licensure Primary Study MICP11: Primary Wheezing Safety Finding: Medically Significant Wheeze (MSW) FluMist® vs. TIV

- Age 6-23 months: Increased risk for MSW after FluMist®
- Age 24-59 months: No increased risk for MSW after FluMist®

Age Mo.	FluMist® N (%)*	TIV N (%)*	Relative Risk (95% CI)
6-23	117 (5.9%)	75 (3.8%)	1.55 (1.17, 2.05)
6-11 [†]	47 (6.9%)	29 (4.2%)	1.62 (1.04, 2.53)
12-23 [†]	70 (5.4%)	46 (3.6%)	1.50 (1.05, 2.16)
24-59	46 (2.1%)	56 (2.5%)	0.83 (0.56, 1.21)

*Number in FluMist®=4179; TIV=4173.

[†]Not a pre-specified age group for analysis

Sources: FDA and MedImmune briefing documents, FDA VRPBAC meeting, May 16, 2007; Belshe NEJM 2007.



Pre-licensure Primary Study MICP11: Reactogenicity and Serious Adverse Events

- Compared with TIV recipients, FluMist® recipients aged 6-59 months had
 - More frequent reports of runny nose/ nasal congestion and low-grade fever after 1st dose
 - Similar serious adverse event (SAE) rates through 180 days (3.3 % FluMist® vs. 3.1% TIV)
- In a post-hoc analysis, Flumist® recipients aged 6-11 months had increased risk for all-cause hospitalization through 180 days:
 - FluMist® (6.1%) vs. TIV (2.6%)



Sources: FDA and MedImmune briefing material, FDA VRPBAC meeting, May 16, 2007; Belshe NEJM 2007.

Adverse Events after FluMist®

Reported to VAERS

8/1/2003–2/28/2007


Age Group	All Reports	Serious Reports (%)*
6-23 months*	1	0
2-4 years*	14	0
5-17 years	266	14 (5.2%)
18-49 years	370	27 (7.3%)
≥50 years*	26	1 (3.8)
Unknown	67	1 (1.5%)
Total	744‡	43 (5.8%)

* Percentage of total reports in corresponding age-group

† Outside the licensed age group

‡ Crude reporting rate 10.9 total reports per 100,000 doses distributed

Source: Hua W, Haber P, Izurieta H, unpublished VAERS data



VAERS FluMist® Reports: Selected Respiratory Conditions 8/1/2003–2/28/2007

Age	All reports	Reported main condition (%)*		
		Asthma/ wheezing	Pneumonia	Bronchitis
6-23 mo	1	0	0	0
2-4 yrs	14	0	0	0
5-17 yrs	266	16 (6%)	7 (3 %)	2 (1 %)
≥18 yrs	396	5 (13%)	6 (2%)	4 (1%)
Unknown	67	1 (2%)	2 (3%)	2 (3%)
Total	744	22 (3%)	15 (2%)	8 (1%)

* Percentage of total reports in corresponding age-group
Source: Hua W, Haber P, Izurieta H, unpublished VAERS data



VAERS FluMist® Reports of Asthma/Wheeze by Medical History 8/1/2003–2/28/2007

Age	All	Asthma/Wheezing Adverse Event Report			
		Total*	Asthma History/ Wheeze	Other chronic condition history	No history
6-23 mo	1	0	0	0	0
2-4 years	14	0	0	0	0
5-17 years	266	16 (6%)	7	0	9
≥18 years	396	5 (13%)	3	0	2
Unknown	67	1 (2%)	0	0	1
Total	744	22 (3%)	10	0	12

* Percentage of total reports in corresponding age-group
Source: Hua W, Haber P, Izurieta H, unpublished VAERS data



Key Safety Questions about FluMist®

Use in Young Children

- Will screening for a history of wheezing identify children appropriately?
- Is the safety profile of FluMist® within 6 weeks after vaccination comparable to TIV in children aged <5 years?
 - Is the risk for medically-attended wheeze increased?
 - Is the risk for all-cause hospitalization increased?
 - Are there other unexpected safety signals?
- Are children with a history of wheezing at increased risk for wheeze or hospitalization after FluMist®, compared with those with no history?

VAERS Post-Marketing Surveillance for FluMist® Vaccine

- National passive surveillance system for reporting vaccine adverse events
- Limitations
 - Risk of underreporting or reporting bias
 - Possibly incomplete, unreliable data
- CDC / FDA scientists will review daily alerts of serious adverse event (SAE) reports* and other medically important conditions, including wheezing after FluMist®

*Code of Federal Regulations 21CFR600.80 Serious adverse event defined as involving hospitalization or prolongation of hospitalization, death, life-threatening illness or permanent disability



Vaccine Safety Datalink Project (VSD)

- Collaboration between CDC and 8 managed care organizations MCOs with comprehensive medical and immunization histories of 5.5 million people per year
- Limitations
 - Limited power to assess risk for rare adverse events
 - Inability to monitor vaccine safety in near real-time if vaccine uptake is limited in MCOs

Vaccine Safety Datalink (VSD) and FluMist®

- To date, limited use of FluMist® in VSD sites, except as part of ongoing Phase 4 study*
- VSD will examine rates of clinically important outcomes in children aged <5 years, including
 - Medically-attended wheezing
 - All-cause hospitalization
- FDA licensure and ACIP recommendation decisions and uptake in VSD sites will affect specific aims and methods of VSD study/studies

*CDC has no involvement with this study; sources: MedImmune briefing document, FDA VRPBAC meeting, May 16, 2007 and Product Approval Information 2003, available at <http://www.fda.gov/cber/approvltr/inflmed061703L.htm>



Additional FluMist® Safety Activities

- The Brighton Collaboration to develop standardized case definitions for wheezing/asthma outcomes
- Education and outreach to providers and parents
 - Vaccine Information Statement
- MedImmune phase 4 study*
 - Population of 60,000 FluMist® recipients aged 5-49 years within the Kaiser Permanente system
 - Results anticipated 2011

*CDC has no involvement with this study; sources: MedImmune briefing document, FDA VRPBAC meeting, May 16, 2007 and Product Approval Information 2003, available at <http://www.fda.gov/cber/approvltr/inflmed061703L.htm>



Acknowledgements

FDA

- Hector Izurieta
- Wae Hua
- Sang Ahnn

Kaiser Permanente CO

- Matt Dailey
- Simon Hambidge
- Jason Glanz

NVPO

- Ben Schwartz

CDC

- Carolyn Bridges
- James Baggs
- Tony Fiore
- Paul Gargiullo
- Julianne Gee
- Jane Gidudu
- Penina Haber
- John Iskander
- Elaine Miller
- Miriam Sabin
- David Shay
- Eric Weintraub
- Claudia Vellozzi



Background Slides



FluMist® Product History

- **June 2003:** FluMist (frozen) licensed for use in healthy persons aged 5-49 years (should not be administered to individuals without a history of asthma or reactive airway disease)
- **January 2007:** FluMist liquid formulation licensed to replace frozen formulation (first widespread use anticipated during 2007/08 influenza season)
- **May/June 2007:** FDA Vaccine and Related Product Biologics Advisory Committee (VRPBAC) reviewed MedImmune's Biologics License Application supplement for use of FluMist (liquid) in children aged 12-59 months without a history of wheeze/asthma; currently under FDA review



FDA Vaccines and Related Product Advisory Committee Meeting, May 16, 2007

Questions 2: Do the data demonstrate that the benefits will exceed the risk of FluMist® for use in

- A. The applicant's proposed population (children 12 to 59 months of age without a history of wheeze/asthma)
 - Vote –Yes: 9 and No: 6
- B. Children in the age strata 6-23, regardless of wheezing history?
 - Vote –Yes: 3 and No: 12
- C. Children in the age strata 24-59 months, regardless of wheezing history?
 - Vote –Yes: 15 and No: 0

Source VRPBAC transcript, page 256. available at
<http://www.fda.gov/ohrms/dockets/ac/07/transcripts/2007-4292t1.pdf>



Pre-licensure Primary Study MICP11: Wheezing Outcomes*

- Primary: Medically Significant Wheeze (MSW):
Wheezing on physical exam and ≥ 1
 - Sign of respiratory distress (tachypnea, retractions, or dyspnea)
 - Hypoxemia (O2 sat $< 95\%$), or
 - New prescription for daily bronchodilator therapy (not on an “as needed” basis)
- Secondary: “Any wheezing” reported by clinician or parent
 - MedDRA preferred terms: wheeze, asthma, bronchospasm, bronchilolitis

*Occurring from receipt of dose 1 through 42 days after the last dose
Sources: FDA and MedImmune briefing documents, FDA VRPBAC meeting, May 16, 2007; Belshe NEJM 2007.



Use of Wheezing History and Outcomes in FluMist® Study MICP11

- Exclusion Criteria: Severe asthma, wheeze within 42 days
- Criteria for stratification: recurrent wheeze history
- Post-hoc analysis to assess association between history of wheeze and certain adverse events
- Pre-specified safety outcomes
 - Primary: Medically significant wheeze (MSW)
 - Secondary: Any wheezing



Selected Adverse Events after FluMist® Reported to VAERS: 8/1/2003–7/31/2005

Main Condition	All Reports (%)*	Serious Reports (%)*
Respiratory	217 (47.2%)	15 (37.5%)
Asthma	12	5
Pneumonia	10	5
Constitutional	67 (14.6%)	4 (10%)
Allergic	54 (11.7%)	6 (15%)
Abdominal	33 (7.2%)	1 (2.5%)
Ear-nose-throat	18 (3.9%)	0
Neurological	10 (2.3%)	7 (17.5%)
GBS*	3	3
Bell's Palsy	1	1
Cardiovascular	10 (2.2%)	3 (7.5%)
Ocular	7 (1.5%)	1 (2.5%)
Other	44 (9.6%)	3 (7.5%)
Total	460 (100%)	40 (100%)

*Guillain-Barré Syndrome

Source: Izurieta, Haber. *JAMA*. 2005.



Adverse Events after FluMist® in 15 Children Aged <5 Years: Main Condition Reported to VAERS 8/1/2003–2/28/2007

Adverse Event	N
Vaccine administration error	6
Secondary transmission	2
Rash/ other allergic	2
Rhinitis	2
Abdominal	1
Seizure	1
Vaccine failure	1

Source: Hua W., Haber P, Izurieta H, unpublished VAERS data

